

# 15 General Pest Management Considerations – Apricots

## 15.1 Diseases

### Bacterial Canker (*Pseudomonas syringae*)

#### • Biology & Cultural

See the description of pathogen biology and copper spray timing for this disease under “Cherries—Diseases.” No recent research has been done on bacterial canker in apricots under east coast conditions. However, this pathogen is probably responsible for much of the early tree death that follows limb die-back and trunk cankers that commonly develop in young apricot plantings. *P. syringae* can also cause a severe blossom blast on apricots if pathogen populations are high when a bloom-time frost occurs. Blossom blast can kill most of the spurs on affected trees.

To minimize the potential for severe losses from bacterial canker, apricot plantings should not be located adjacent to old sweet cherry orchards nor close to hedgerows or woodlots that contain wild *Prunus* (especially wild black cherry, *Prunus serotina*). Copper sprays should be applied in spring and fall as for sweet cherries. The spring copper spray may be even more important on apricot than on sweet cherry because apricots appear more sensitive to blossom blast. Entry of the pathogen through pruning wounds can be minimized if pruning is done during hot, dry weather after bloom or after harvest, because *P. syringae* is suppressed by hot weather. Trunks and scaffolds on apricot trees should be painted with white latex paint in autumn to reduce the potential for winter injury, as *P. syringae* can also enter where bark is injured during winter.

#### • Pesticide Application Notes

[1.1] Optimum timing and effectiveness of copper applications for control of bacterial canker and blossom blast of apricots has not been determined under eastern conditions. However, applying copper at the maximum labeled rates at the late dormant stage should increase the likelihood that effective copper residues will remain on the trees throughout the bloom period. Copper applications are especially important on young trees because trees less than five years old can be killed by bacterial canker if infections reach the main trunk.

### Bacterial Spot (*Xanthomonas arboricola* pv. *pruni*)

#### • Biology & Cultural

Bacterial spot can be devastating to apricots. Apricot varieties developed in drier climates and then grown in the more humid climate of NY are the most likely to be susceptible. This disease will be more severe in the warmer southern portions of NY, in wet years, in orchards with lighter (sandy) soils, and in windy orchard sites. The bacterial spot pathogen, *Xanthomonas arboricola* pv. *pruni* infects leaf scars at leaf drop and overwinters in infected twigs. Bacteria populations subsequently multiply during

warm weather and ooze out during spring rains. Immature tissues are less susceptible to the bacterial infection, and as such, infections will not begin until petal fall/shuck split. Early season copper applications applied to manage bacterial canker are quite effective for controlling the bacterial spot populations, but also likely to induce phytotoxicity if one is not careful.

#### • Pesticide Application Notes

Unfortunately, there are no materials registered for bacterial spot on apricots. Despite the effectiveness, do not make a dormant copper application for bacterial spot. Copper applications to manage bacterial canker and bacterial blast are still allowed whether or not the planting has bacterial spot.

### Brown Rot

#### • Biology & Cultural

Refer to the reference materials list at the end of this publication for a Fact Sheet containing details on the biology and management of this disease.

#### • Pesticide Application Notes

[2.1] Apricots are much more susceptible to blossom blight than any other stone-fruit species. At least 1 protective spray should be applied each year; repeat at full bloom and/or petal fall if wet weather prevails during bloom. Pristine, Indar, and Orbit are generally more effective than captan or Bravo. When used at a rate of 10 oz/100 gal dilute, Rovral provides 24–48 hr kickback activity against blossom blight infections at 68° F; Indar and Orbit also have significant kickback activity. Scala, Vanguard, and Elevate have not been tested on apricots, but they have been effective for blossom blight on peaches.

[2.2] The shuck split application is an important spray. Chlorothalonil (Bravo, Echo) provides a longer period of protection than either captan or sulfur, but do not use Chlorothalonil after shuck split.

[2.3] Fruit are very susceptible to brown rot prior to pit hardening and the last 3 wk before harvest. It is recommended that spray intervals be shortened during the preharvest period and that Indar, Orbit, or Pristine be used if disease pressure is high (warm, wet). Indar and Orbit are registered for use beginning 3 weeks before harvest. Of these materials, Indar has the longest residual activity.

#### • Pesticide Resistance

[2.4] For resistance management purposes, the SI fungicides (Indar, Orbit) should not be used routinely throughout the season for BOTH blossom blight AND fruit rot control.

## Peach Scab

### • Biology & Cultural

Peach scab can severely damage apricot if spring weather is warm and wet and no fungicides are applied at shuck split and first cover. The disease is more common following a year when spring frosts caused a crop failure, because trees grown for an entire summer without fungicides are more likely to carry peach scab infections the following year.

### • Pesticide Application Notes

[3.1] Apply 2 or 3 sprays at 10–14-day intervals beginning at shuck split. Under light disease pressure, a single application of Bravo or Echo applied at shuck split may provide season-long control. Bravo and Echo cannot be applied after shuck split.

## Perennial (Cytospora, Valsa) Canker

### • Biology & Cultural

[4.1] See discussion of this disease under Peaches. Also, refer to the reference materials list at the end of this publication for a Fact Sheet containing details on the biology and management of this disease.

## Phytophthora Root, Crown, and Collar Rots

### • Biology & Cultural

[5.1] Apricot rootstocks are perhaps more susceptible to Phytophthora root, crown, and collar rots than any other fruit tree rootstock grown in New England. The main defenses against these diseases should be providing good soil drainage through proper site selection and physical manipulations such as tiling or planting on berms. In general, berms are much more effective than tiling. See comment [5.2] about pesticide applications.

Refer to the reference materials list at the end of this publication for a Fact Sheet containing details on the biology and management of this disease.

### • Pesticide Application Notes

[5.2] In addition to the cultural practices noted in comment [5.1], Ridomil will provide additional protection in wet years, on marginal sites, or in wetter sections of the orchard. Applications should be made just before growth starts in the spring and at 2–3 month intervals thereafter if soil conditions are very wet. Apply to the soil beneath the tree canopy in sufficient water to assure good coverage (material is moved into the soil by subsequent rain or irrigation). See label for further details.

## 15.2 Insects and Mites

### European Red Mite, Twospotted Spider Mite

#### • Biology & Cultural

Refer to the reference materials list at the end of this publication for a Fact Sheet containing details on the biology and management of this pest.

#### • Pesticide Application Notes

[6.1] Apply as mites appear in a minimum of 50 gal/A. Limited to 1 application per season.

[6.2] Use lower rate of Nexter for European red mite, and higher rate of Nexter, for twospotted spider mite.

[6.3] Non-bearing trees only.

## Oriental Fruit Moth

### • Biology & Cultural

Refer to the reference materials list at the end of this publication for a Fact Sheet containing details on the biology and management of this pest.

### • Biological & Non-chemical Control

[7.1] Pheromone disruption is economically justified if 2–3 sprays are normally applied, and if no other insecticide sprays are routinely needed for other pests after petal fall. For this reason, disruption may not be economical for the 1st brood, as plum curculio sprays at this time normally would also control oriental fruit moth. Pheromones should be applied in mid-June before initiation of the 2nd flight; the need for re-application depends on residual field life of specific formulations: Isomate-M 100, 90 days; Checkmate and 3M SprayableOFM-F, 14 days. The residual life of insecticide sprays or a double rate of the 3M sprayable deposit can be extended by the addition of pheromone. Pheromone may be needed in border rows of a spreader-sticker such as Nu-Film-17 at 1 pt/A. Border insecticide sprays may be needed in orchards adjacent to sources of adult immigration or in other high pressure situations.

### • Pesticide Application Notes

[7.2] Summer sprays should be timed to start approximately at the 10% hatch point, 175–200 DD (base 45° F) after the first adult catch of the second brood, with a second application in 10–14 days. In high pressure blocks, a final spray should be applied 2 wk before harvest to control late season larvae. Suggested action threshold: Avg. of >10 adults/ week caught per pheromone trap.

## Peachtree Borers (including Lesser peachtree borer)

### • Biology & Cultural

Refer to the reference materials list at the end of this publication for a Fact Sheet containing details on the biology and management of this pest.

### • Biological & Non-chemical Control

[8.1] Hang pheromone traps at 100/acre in late May before lesser peachtree borer flight begins; use 200/acre if population is predominantly peachtree borer.

### • Pesticide Application Notes

[8.2] Up to 3 sprays of pyrethroids to trunk and scaffold limbs against larvae: June 1–10, July 7–15, and August 1–10. OR 1 application of \*Thionex immediately

after harvest; do not spray fruit. \*Baythroid not labeled for peachtree borer. Suggested action threshold: 1st emergence of adults plus 8 days (in blocks with a history of damage), or 1-2 larvae/tree.

## Plum Curculio

### • Biology & Cultural

Refer to the reference materials list at the end of this publication for a Fact Sheet containing details on the biology and management of this pest.

### • Pesticide Application Notes

[9.1] 2–3 applications. Begin at fruit set (shucks on) and follow at 8–10-day (for Sevin) or 10–14-day (for other products) intervals.

[9.2] Frequent applications (7–10-day intervals) of Surround and maximal coverage (minimum of 100 gal/A) are advised while there is active foliar growth.

## Tarnished Plant Bug

### • Biology & Cultural

Refer to the reference materials list at the end of this publication for a Fact Sheet containing details on the biology and management of this pest. Satisfactory control requires adequate management of orchard weeds that attract this pest and act as alternate hosts.

### • Monitoring & Forecasting

[10.1] Most catfacing injury is caused before shuck split. Later season feeding generally results in only minor surface scarring.

[10.2] Apply spray as insects or damage appears. Suggested action threshold: 3 bleeding sites/tree.

### • Pesticide Application Notes

[10.3] At 10–14-day intervals as needed in midsummer. Suggested action threshold: 3 bleeding sites/tree.

## Western Flower Thrips

### • Biology & Cultural

Drought conditions and high temperatures may encourage damaging populations. Adults move from alternate weed or crop hosts to fruit just prior to and during harvest, feed on the fruit surface in protected sites, such as in the stem end, the suture, under leaves and branches, and between fruit. Feeding results in silver stipling or patches; injury is particularly obvious on highly colored varieties.

### • Pesticide Application Notes

[11.1] In orchards with severe infestations, a petal fall application may be warranted against thrips feeding in fruit clusters. Control may be improved by addition of an adjuvant.

[11.2] An application immediately after harvest may prevent subsequent losses; however, an additional application may be needed if pressure is severe. Note PHI. Control may be improved by addition of an adjuvant.

## Storage Rots

### • Pesticide Application Notes

[12.1] A postharvest treatment with Scholar SC via dipping, flooders, T-jet, or similar system for control of storage rots is recommended for fruit coming from orchards where sporulating brown rot was observed, or when one hopes keep fruit in cold storage for a few days prior to sale. Holding tanks in postharvest treatment equipment must have excellent agitation to keep fungicides in suspension. Solutions must be replenished regularly as directed on the product label. Never exposed treated fruit to direct sunlight. This will cause the fungicide to break down.

### 15.3 Apricot Spray Tables

**Table 15.3.1. Pesticide Spray Table -- Apricots.**

Refer to inside back cover for key to abbreviations and footnotes.

Pest	Product	Rate	REI (hrs)	PHI (days)	Comments (see text)
<b>Late Dormant</b>					
<b>Phytophthora root, crown and collar rots</b>	Ridomil Gold 4EC	1.5 fl oz/1,000 sq ft treated	12	0	[5.1]
<b>Bacterial canker (<i>Pseudomonas syringae</i>)</b>	Kocide 40DF	2-4 lb/100 gal (max 12 lb/A)	24	BL,PH(C)	[1.1]
	or Kocide 50WP or §Cuprofix	10-16lb/A	24	BL,PH(C)	
	Disperss 40DF or other coppers	See comments			
<b>Popcorn</b>					
<b>Brown rot (Blossom blight)</b>	Bravo Ultrex 82.5WDG or Bravo Weather Stik 6F or other chlorothalonil formulations (see labels)	0.9-1.25 lb/100 gal 1.0-1.4 pt/100 gal	12hr/7days(E)	SS	
	Captan 50WP or Captan 4L	2 lb/100 gal(max 5lb/A) 1 qt/100 gal (max 5 qt/A)	96 (E) 24 (E)	0	[2.1]
	<i>OR</i> Echo 6F or Echo 90DF	1.0-1.4 pt/100 gal 0.75-1.2 lb/100 gal	12 hr/7days(E)	SS	
	<i>OR</i> Elevate 50WDG	0.33-0.5 lb/100 gal (max 6 lb/A)	12	0	
	<i>OR</i> Indar 75WS	0.8 oz/100 gal (max 2 oz/A)	12	0	
	<i>OR</i> Orbit 3.6EC	1.6 fl oz/100 gal (max 4 fl oz/A)	24	0	
	<i>OR</i> Rovral 50WP or Rovral 4F	8-10 oz/100 gal (max 2 lb/A) 8-10 fl oz/100 gal	24	PF	
	<i>OR</i> Pristine 38WDG	10.5-14.5 oz/A	12	0	
	<i>OR</i> Scala 600SC	9-18 fl oz/A	12	2	
	<i>OR</i> Vanguard 75WG	5 oz/A	12	BL	
<b>Tarnished plant bug</b>	See materials listed under Petal Fall.				[10.1]
<b>Bloom</b>					
<b>Brown rot (blossom blight)</b>	See materials and comments listed under Popcorn.				
<b>Petal Fall</b>					
<b>Brown rot (Blossom blight)</b>	See materials and comments listed under Popcorn, except Vanguard, which cannot be used after bloom.				
<b>Plum curculio</b>	Assail 30SG	5.3-8 oz/Acre	12	7	
	*Imidan 70WP	3/4-1 lb/100 gal	72	14	[9.1]
	<i>OR</i> *Baythroid 2E or *Baythroid XL 1L	2.4-2.8 fl oz/A 2.4-2.8 fl oz/A	12 12	7 7	
	<i>OR</i> Sevin XLR Plus, 4F or Sevin 80, *80WS	2-3 qt/A 2.5-3.75 lb/A	12	3	
	<i>OR</i> §Surround 95WP	50 lb/100 gal	4	0	[9.2]

**Table 15.3.1. Pesticide Spray Table -- Apricots.**

Refer to inside back cover for key to abbreviations and footnotes.

<b>Pest</b>	<b>Product</b>	<b>Rate</b>	<b>REI (hrs)</b>	<b>PHI (days)</b>	<b>Comments (see text)</b>
<b>Peachtree borers (including Lesser peachtree borer)</b>	Pheromone disruption ties: §Isomate-LPTB				[8.1]
<b>Tarnished plant bug</b>	*Asana XL 0.66EC	2-5.8 oz/100 gal	12	14	[10.2]
	<i>OR</i> Assail 30SG	5.3-8 oz/Acre	12	7	
	<i>OR</i> *Baythroid 2E	2.0-2.4 fl oz/A	12	7	
	or *Baythroid XL 1L	2.0-2.4 fl oz/A	12	7	
	*Proaxis 0.5CS	2.6-5.1 fl oz/A	24	14	
	<i>OR</i> *Warrior 1CS	2.6-5.1 fl oz/A	24	14	
<b>Western flower thrips</b>	SpinTor 2SC	4-8 fl oz/A	4	14	[20.2]
	or §Entrust 80WP	1.25-2.5 oz/A			
<b>Shuck Split</b>					
<b>Brown rot (Blossom blight)</b>	Bravo Ultrex 82.5WDG	0.9-1.25 lb/100 gal	12hr/ 7days(E)	SS	
	or Bravo Weather Stik 6F	1.0-1.4 pt/100 gal			
	or other chlorothalonil formulations (see labels)				
	Captan 50WP	2 lb/100 gal(max 5 lb/A)	96 (E)	0	[2.2]
	or Captan 4L	1 qt/100 gal(max 5 qt/A)	24 (E)		
	<i>OR</i> Echo 6F	1.0-1.4 pt/100 gal	12hr/ 7days(E)	SS	
	or Echo 90DF	0.75-1.2 lb/100 gal			
	<i>OR</i> Pristine 38WDG	10.5-14.5 oz/A)	12	0	
	<i>OR</i> §Sulfur 95WP	5 lb/100 gal	24	0	
<b>Peach Scab</b>	Any of the products listed above for brown rot at shuck split.				[3.1]
	<i>OR</i> Gem 500SC	1.9-3.8 oz/A	12	1	
<b>Additional Summer Sprays</b>					
<b>Brown rot (Blossom blight)</b>	Captan 50WP	2 lb/100 gal (max 5 lb/A)	96 (E)	0	[2.3]
	Captan 4L	1 qt/100 gal (max 5 qt/A)	24 (E)		
	<i>OR</i> Elevate 50WDG	0.33-0.5 lb/100 gal (max 6 lb/A)	12	0	
	<i>OR</i> Indar 75WS	0.8 oz/100 gal (max 2 oz/A)	12	0	
	<i>OR</i> Orbit 3.6EC	1.6 floz/100 gal (max 4 fl oz/A)	24	0	
	<i>OR</i> Pristine 38WDG	10.5-14.5 oz/A)	12	0	
	<i>OR</i> §Sulfur 95WP	5 lb/100 gal	24	0	
<b>European red mite,</b>	Apollo 4SC	2-8 oz/A	12	21	[6.1]
<b>Twospotted spider mite</b>	<i>OR</i> §Nexter 75WS	4.4-10.7 oz/A	12	300(PH)	[6.2]
	<i>OR</i> Savey 50DF	3-6 oz/A	12	28	
<b>Oriental fruit moth</b>	Pheromone disruption ties: §3M Sprayable				[7.1]
	Pheromone for OFM	1.7 oz/A			
	or §Checkmate OFM-F	1.32-2.93 fl oz/A			
	§Isomate-M 100	100 ties/A			
	<i>OR</i> *Asana XL 0.66EC	2-5.8 oz/100 gal	12	14	[7.2]
	<i>OR</i> Assail 30SG	5.3-8 oz/Acre	12	7	

**Table 15.3.1. Pesticide Spray Table -- Apricots.**

Refer to inside back cover for key to abbreviations and footnotes.

Pest	Product	Rate	REI (hrs)	PHI (days)	Comments (see text)
<b>Oriental fruit moth</b> (continued)	OR *Baythroid 2E	2.0-2.4 fl oz/A	12	7	
	or *Baythroid XL 1L	2.0-2.4 fl oz/A	12	7	
	OR *Proaxis 0.5CS	2.6-5.1 fl oz/A	24	14	
	OR Sevin XLR Plus, 4F or Sevin 80S, *80WS	2-3 qt/A 2.5-3.75 lb/A	12	3	
	OR *Warrior 1CS	2.6-5.1 fl oz/A	24	14	
<b>Peachtree borers</b> (including Lesser peachtree borer)	*Asana XL 0.66EC	2-5.8 oz/100 gal	12	14	[8.2]
	OR *Baythroid 2E or *Baythroid XL 1L	1.4-2.0 fl oz/A 1.4-2.0 fl oz/A	12 12	7 7	
	OR *Proaxis 0.5CS	2.6-5.1 fl oz/A	24	14	
	OR *Thionex 3EC or *Thionex 50WP	1 qt/100 gal 1.5 lb/100 gal	24 21	21/30(A)	
	OR *Warrior 1CS	2.6-5.1 fl oz/A	24	14	
<b>Tarnished plant bug</b> <b>Stink bug</b>	*Asana XL 0.66EC	2-5.8 oz/100 gal	12	14	[10.3]
	OR Assail 30SG	5.3-8 oz/Acre	12	7	
	OR *Baythroid 2E or *Baythroid XL 1L	2.0-2.4 fl oz/A 2.0-2.4 fl oz/A	12 12	7 7	
	OR *Warrior 1CS	2.6-5.1 fl oz/A	24	14	
<b>Western flower thrips</b>	SpinTor 2SC or §Entrust 80WP	4-8 fl oz/A 1.25-2.5 oz/A	4	14	[11.2]
<b>Postharvest</b>					
<b>European red mite, Twospotted spider mite</b>	Nexter 75WS	4.4-10.7 oz/A	12	300(PH)	[6.2]
<b>Peachtree borers</b>	*Thionex 50WP	1.5 lb/100 gal	24	21	[8.2]
	or *Thionex 3EC	1 qt/100 gal	30		
<b>Autumn</b>					
<b>Bacterial canker</b> ( <i>Pseudomonas syringae</i> )	Kocide 40DF	2-4 lb/100 gal (max 12 lb/A)	24	BL,PH(C)	[1.1]
	or Kocide 50WP or §Cuprofix Disperss 40DF	10-16 lb/A	24	BL,PH(C)	
	or other coppers	See comments			

**Table 15.3.2. Plant Growth Regulator Use in Apricots**

Refer to inside back cover for key to abbreviations and footnotes.

Timing	Product	Concentration	Rate of Formulated Product	Comments
<b>PREHARVEST FRUIT DROP CONTROL</b>				
1-2 weeks before anticipated harvest	ReTain	132 ppm	0.74 lb/acre or 333 g/acre (1 pouch)	Apply in sufficient water to ensure thorough but not excessive coverage. An organosilicone surfactant (12 oz/100 gal) should be used with ReTain.